

Providing an improved wind stress parameterization and QuikSCAT wind stress constraints for ECCO2

David Moroni₁
Holger Brix₂
Dimitris Menemenlis₁
Hong Zhang₂

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



www.nasa.gov

1. Jet Propulsion Laboratory / California Institute of Technology
2. University of California, Los Angeles

11/10/2009



ECCO2 Meeting – CalTech, Pasadena, CA

1





Motivation

- More than 10 years of QuikSCAT data are available at 90% global daily coverage provides exciting opportunity for improving the ECCO2 MITgcm.
- An improved, 'prototype' wind stress analysis product would have enormous impacts for both operations and research.
- Potential for optimizing of high resolution simulations.
- Potential contributions to the fundamental understanding of the physics and dynamics of oceanographic processes.






Objectives

1. Test an improved wind stress parameterization in MITgcm.
 - Run surface-relative and absolute wind integrations.
 - Compare with Large and Pond (1981) parameterization.
 - Compare all runs with QuikSCAT retrievals.
2. Incorporate improved parameterization into MITgcm.
3. Quantify model vs. data errors through comparisons of QuikSCAT-ECCO2 wind stress.
 - Requirement for assimilation.
4. Repeat $1/8^\circ$ simulation (Hill et al. 2007) using both:
 - QuikSCAT wind stress constraints and
 - improved, sea-state-dependent (hypothetically ideal) surface wind stress parameterization.



Power Spectrum: MITgcm vs. TOPEX







Relative vs. Absolute Wind



Mean Surface Velocities: 1992-2002

Zonal: Absolute – Relative




Meridional: Absolute – Relative







Latent Heat Flux: Global Probability Density





Momentum Flux: Global Probability Density







Comparisons to In Situ Stress


- Recent developments in sea-state dependent stress parameterizations have been adjusted to provide a better fit to in situ stress observations (Bourassa 2004, 2006):
- Created a 'prototype' QuikSCAT Level-2 wind stress retrieval product based on the Bourassa (2006) parameterization.







QuikSCAT-ECCO2 Inconsistencies

Zonal Stress RMS Difference (N m^{-2})




> 95% Data Errors Below 0.025 N m^{-2}



Concluding Remarks

- An improved $1/8^\circ$ simulation would raise the bar for higher resolution simulations.
- Global, gridded stress fields from ECCO2 could be provided as an open-source analysis product.
 - Potential for climate studies and short-term forecasting applications.
 - Potential for cal/val with other OGCMs.
 - Potential for cal/val with current and future scatterometer stress GMF development.
- Re-estimation of wind work upon the ocean circulation, continuing in the direction of Wunsch (1998) and von Storch et al. (2007).



Thank you!